

survival factor in late-stage melanoma cells, p38 attenuates Fas expression via inhibition of NF- κ B. The present invention is based, in part, on an investigation of whether ATF2-derived peptides could be used to alter the sensitivity of human melanoma cells to radiation and chemical treatment. Of four 50 amino acid peptides tested, the peptide spanning amino acid residues 50-100 elicits the most efficient increase in the sensitivity of human melanoma cells to UV radiation or treatment by mitomycin C, Adriamycin[®] ~~adriamycin~~ and verapamil, or UCN-01, as revealed by apoptosis assays. Sensitization by ATF2 peptide was also observed in the MCF7 human breast cancer cells, but not in early-stage melanoma, or melanocytes, or in *in vitro* transformed 293T cells. When combined with an inhibitor of the p38 catalytic activity, cells expressing the 50-100 fragment of ATF2 exhibited an increase in the degree of programmed cell death, indicating that combined targeting of ATF2 and p38 kinases is sufficient to induce apoptosis in late-stage melanoma cells. The peptide's ability to increase levels of apoptosis coincided with increased cell surface expression of Fas, which is the primary death-signaling cascade in these late stage melanoma cells. Overall, our studies identify a critical domain of ATF2, which may be used to sensitize tumor cells to radiation and chemical treatment-induced apoptosis and which can induce apoptosis, when combined with inhibition of ATF2 kinase, p38.

Please replace the paragraph at page 9, line 29 to page 10, lines 6 with the following:

“Inhibition of ATF2 activity” (and all grammatical variations thereof) includes, but is not limited to, inhibition of ATF2-regulated transcription; inhibition of tumor cell growth (relative to untreated tumor cells); an increase in apoptosis; an increase in the sensitivity of tumor cells, particularly human melanoma and breast cancer cells, to UV radiation or treatment by chemotherapeutic drugs such as mitomycin C, Adriamycin[®] ~~adriamycin~~ and verapamil, and UCN-01; and the like. In particular, inhibition of ATF2 activity comprises inhibiting growth of a tumor cell, which method comprises inhibiting transcriptional activity of ATF2.

Please replace the paragraph at page 16, line 25 to page 17, line 21 with the following:

Vectors typically comprise the DNA of a transmissible agent, into which foreign DNA is inserted. A common way to insert one segment of DNA into another segment of DNA involves the use of enzymes called restriction enzymes that cleave DNA at specific sites (specific groups of nucleotides) called restriction sites. A "cassette" refers to a DNA coding sequence or segment of DNA that codes for an expression product that can be inserted into a vector at defined restriction sites. The cassette restriction sites are designed to ensure insertion of the cassette in the proper reading frame. Generally, foreign DNA is inserted at one or more restriction sites of the vector DNA, and then is carried by the vector into a host cell along with the transmissible vector DNA. A segment or sequence of DNA having inserted or added DNA, such as an expression vector, can also be called a "DNA construct." A common type of vector is a "plasmid", which generally is a self-contained molecule of double-stranded DNA, usually of bacterial origin, that can readily accept additional (foreign) DNA and which can readily introduced into a suitable host cell. A plasmid vector often contains coding DNA and promoter DNA and has one or more restriction sites suitable for inserting foreign DNA. Coding DNA is a DNA sequence that encodes a particular amino acid sequence for a particular protein or enzyme. Promoter DNA is a DNA sequence which initiates, regulates, or otherwise mediates or controls the expression of the coding DNA. Promoter DNA and coding DNA may be from the same gene or from different genes, and may be from the same or different organisms. A large number of vectors, including plasmid and fungal vectors, have been described for replication and/or expression in a variety of eukaryotic and prokaryotic hosts. Non-limiting examples include pKK plasmids (Clontech® Clontech), pUC plasmids, pET plasmids (Novagen, Inc., Madison, WI), pRSET or pREP plasmids (Invitrogen, San Diego, CA), or pMAL plasmids (New England Biolabs, Beverly, MA), and many appropriate host cells, using methods disclosed or cited herein or otherwise known to those skilled in the relevant art. Recombinant cloning vectors will often

include one or more replication systems for cloning or expression, one or more markers for selection in the host, *e.g.* antibiotic resistance, and one or more expression cassettes.

Please replace the paragraph at page 25, line 23 to page 26, line 1 with the following:

Various companies produce viral vectors commercially, including but by no means limited to ~~Avigen, Inc.~~ Avigen, Inc.[®] (Alameda, CA; AAV vectors), ~~Cell Genesys~~ Cell Genesys[®] (Foster City, CA; retroviral, adenoviral, AAV vectors, and lentiviral vectors), ~~Clontech~~ Clontech[®] (retroviral and baculoviral vectors), Genovo, Inc. (Sharon Hill, PA; adenoviral and AAV vectors), Genvec (adenoviral vectors), IntroGene (Leiden, Netherlands; adenoviral vectors), Molecular Medicine (retroviral, adenoviral, AAV, and herpes viral vectors), Norgen (adenoviral vectors), Oxford BioMedica (Oxford, United Kingdom; lentiviral vectors), and ~~Transgene~~ Transgene[™] (Strasbourg, France; adenoviral, vaccinia, retroviral, and lentiviral vectors).

Please replace the paragraph at page 28, line 25 to page 29, line 11 with the following:

The therapeutic compositions of the invention can be used in combination with other anti-cancer strategies, as disclosed herein. In particular, as noted above, a particular advantage of ATF2 inhibition in accordance with the invention results from the adjuvant effect of this strategy on traditional tumor therapies. Although the methods of the invention are effective in inhibiting tumor growth and metastasis, the vectors and methods of the present invention are advantageously used with other treatment modalities, including without limitation radiation and chemotherapy. In particular, ATF2 inhibition can be administered with a chemotherapeutic such as, though not limited to, a p38/JAK kinase inhibitor, *e.g.*, SB203580; a phosphatidyl inositol-3 kinase (PI3K) inhibitor, *e.g.*, LY294002; a MAPK inhibitor, *e.g.*, PD98059; a JAK inhibitor, *e.g.*, AG490; preferred chemotherapeutics such as UCN-01, NCS, mitomycin C (MMC), NCS, and anisomycin; taxoids such as Taxol® ~~taxol~~, Taxotere® ~~taxotere~~ and other taxoids (*e.g.*, as

Test compounds are screened from large libraries of synthetic or natural compounds. Numerous means are currently used for random and directed synthesis of saccharide, peptide, and nucleic acid based compounds. Synthetic compound libraries are commercially available from ~~Maybridge Chemical Co.~~ Maybridge Chemical Co.[®] (Trevillet, Cornwall, UK), ~~Comgenex~~ Comgenex[®] (Princeton, NJ), Brandon Associates (Merrimack, NH), and ~~Microsource~~ Microsource[®] (New Milford, CT). A rare chemical library is available from ~~Aldrich~~ Aldrich[®] (Milwaukee, WI). Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available from *e.g.* Pan Laboratories (Bothell, WA) or MycoSearch (NC), or are readily producible. Additionally, natural and synthetically produced libraries and compounds are readily modified through conventional chemical, physical, and biochemical means (Blondelle *et al.*, TIBTech 1996, 14:60).

Cell lines. LU1205 cells (also known as 1205LU), a late-stage human melanoma cell line, were maintained in MCDB130/L15 medium (4:1) supplemented with 5% fetal bovine serum, L-glutamine and antibiotics. LU1205 cells that stably express ATF2-derived peptide II or peptide IV were maintained in the same medium supplemented with G418 (200 mg/ml). The late-stage melanoma cells, FEMX (Ivanov et al., J. Biol. Chem. 1999, 274:14079-14089; and Ivanov et al.,

Oncogene 2000, 19:3003-3012) were maintained in RPMI supplemented with 10% FBS and antibiotics. The medium for WM1552 cells, an early-phase human melanoma cell line, was the same as for the cells (Ivanov et al., Oncogene 2000, 19:933-942), supplemented with insulin (5mg/ml). Human melanocytes (FOM71; Kath et al., Anticancer Res. 1989, 9:865-872) were maintained in MCDB153 (Sigma) medium supplemented with 2% FBS (Cansera), 10% chelated FBS (Sigma), 2 mM L-glutamine (~~Cellgro~~) (Cellgro®), 20 pM cholera toxin (Sigma), 100 nM endothelin 3 (Peninsula), 10 ng/ml stem cell factor (Research & Development) and 250 pM basic fibroblast growth factor (~~GIBCO~~) (GIBCO®).

Please replace the paragraph at page 39, lines 9-16 with the following:

Chemicals. The pharmacological inhibitors of JAKs (AG490), p38 (SB203580) and PI3K (LY294002) were purchased (~~Calbiochem~~) (Calbiochem®). Mitomycin C (MMC), Adriamycin® ~~adriamycin~~ and verapamil ~~Verapamil~~ were purchased from Sigma. The radiomimetic drug neocarzinostatin (NCS) was obtained from Kayaku Co. (Tokyo, Japan). The nuclear export inhibitor Leptomycin B was kind gift of Dr. Yoshida (Kyushu University, Japan) (Kudo et al., Proc. Natl. Acad. Sci. USA 1999, 96:9112-9117). The chemotherapeutic drug 7-hydroxystaurosporine (UCN-01) was kindly provided by the Drug Synthesis and Chemistry Branch at NCI (Gescher, Crit. Rev. Oncol. Hematol. 2000, 34:127-135).

Please replace the paragraph at page 39, lines 17-24 with the following:

Stable transfection and selection. Oligonucleotides corresponding to ATF2 peptides within amino acid residue 1-50 (peptide I), 50-100 (peptide II), 100-150 (peptide III) and 150-200 (peptide IV) were PCR amplified and cloned into *Bam*HI and *Xba*I sites of pcDNA3 (Invitrogen Invitrogen®, Carlsbad, CA), which contains HA-penetratin tag on its NH2-terminal domain. Cloned material was verified via sequencing. pcDNA3-HA-neo or pcDNA3-HA encoding each of

the four peptides was electroporated (230V, 1050 microfarads) into the respective cell lines as previously described (Ronai, Z., et al., Oncogene 1998; 16:523-531). Cells were maintained in G418 (500 μ g/ml) for 2 weeks before mixed population were pooled and characterized.

Please replace the paragraph at page 39, line 25 to page 40, line 6 with the following:

Immunohistochemistry and Western blot analysis. Cells were grown on cover slips before subjected to fixation (3% paraformaldehyde, 2% sucrose in PBS for 10 min at room temperature) followed by permeabilization (0.5% Triton X-100, 3mM MgCl₂, 6% sucrose in PBS for 5 min on ice). Cells were then incubated with antibodies against HA-tag (5 mg/ml) for 1 h at 20°C, before washed with PBS and incubated with secondary (anti-mouse IgG) antibody that is conjugated to FITC (~~Roche Chemicals~~) (Roche® Chemicals) for 1 h at 20°C. Immunofluorescence analysis was carried out using a fluorescence microscope (~~Nikon~~) (Nikon®). Western analysis for the expression of the low molecular weight peptides was carried out using 15% Tricine-SDS-PAGE and antibodies to HA. Secondary antibodies used in this reaction were goat anti mouse IgG conjugated to horseradish peroxidase (1/500). Signals were detected using the ECL system (Amersham-Pharmacia Biotech).

Please replace the paragraph at page 40, lines 7-19 with the following:

Treatment and apoptosis studies of stably transfected melanoma cells. Cells were exposed to UVC at 75J/m² as previously described (Ronai, Z., et al., Oncogene 1998; 16:523-531). SB203580 (1-10 μ M) (~~Calbiochem~~ Calbiochem®, San Diego, CA), NCS (50-100 ng/ml) and mitomycin C (MMC) (0.2-1 μ M) were used to treat melanoma cells. Flow cytometric analysis was performed on a ~~FACS Calibur~~ FACSCalibur™ flow cytometer (Becton Dickinson) using CellQuest™ ~~CellQuest~~ software as described previously (Nicoletti et al., J. Immunol.

Methods 1991; 139:271-279). Cells were pelleted and resuspended in 0.5 ml of hypotonic buffer with 0.1 % Triton X-100 containing PI (40 μ g/ml) and DNase-free RNase A (1 mg/ml). Cells were incubated at 37°C for 30 min and analyzed on a ~~Calibur~~ FACSCalibur™ flow cytometer (Becton Dickinson). The percentage of cells to the left of the diploid G0/1 peak, characteristic of hypodiploid cells that have lost DNA, was defined as the percentage of apoptotic cells. Analysis was performed with light scatter gating. Surface expression of Fas was determined using anti-Fas-PE antibody (~~PharMingen-CA~~) (PharMingen® CA) and flow cytometric analysis. Cell surface expression is measured as mean fluorescence intensity (MFI).

Please replace the paragraph at page 43, lines 2-12 with the following:

Control and peptide expressing cultures were also subjected to treatment with commonly used chemotherapeutic drug Adriamycin® ~~adriamycin~~ alone or in combination with verapamil, which is used to avoid induction of drug resistance. As shown in Figure 2D, sensitivity of LU1205 cells to Adriamycin® ~~adriamycin~~ -induced programmed cell death increased in response to Adriamycin® ~~adriamycin~~ treatment (2-fold when compared with control). Combination of Adriamycin® ~~adriamycin~~ and verapamil caused 4-fold increase in apoptosis of control cells, and an additional (50%) increase in peptide II expressing cells (10-fold increase comparable to neo expressing cells). Peptide IV expressing LU1205 cells exhibited a 70% increase in degree of apoptosis over the control neo expressing cells subjected to the combination of Adriamycin® ~~adriamycin~~ and verapamil (Figure 2D). These observations suggest that the effects mediated by ATF2-peptides are selective to the form of DNA damage and stress.

Please replace the paragraph at page 44, line 28 to page 45, line 16 with the following:

MCF7 is among the better-characterized breast cancer cell lines. The latter resulted in a battery of MCF7-derivatives that were selected for growth based on their ability to develop drug resistance. One such MCF7-derivative is the Adriamycin® ~~adriamycin~~ -resistant MCF7 cell line

(MCF7-Adr), which is 1000 times more resistant to Adriamycin® ~~adriamycin~~ than the original cells line. Peptide II-expressing MCF7-Adr cells exhibited an 8-fold increase in apoptosis following UV-treatment as compared with the 3-fold increase in the MCF7-Adr-neo controls (Figure 4B). This result suggests that peptide II retained its ability to sensitize cells to UV-treatment even if they have acquired resistance to chemotherapeutic drugs. Peptide IV was no longer able to increase the resistance of the cells, as seen in all other cases, but rather sensitized them (from less than 3-fold in controls to greater than 4-fold in MCF7-Adr cells expressing peptide IV). That Adriamycin® ~~adriamycin~~ resistance attenuates peptide IV's ability to elevate resistance of MCF7 cells to UV-treatment implies that Adriamycin® ~~adriamycin~~ and peptide IV may utilize similar cellular pathways required to increase resistance of tumor cells to radiation and chemical treatments. Since they are required to acquire chemotherapeutic resistance, these pathways are no longer affected by peptide IV. The basal degree of apoptosis, which was affected in the MCF7 cells by both ATF2-driven peptides (Figure 4A), was no longer seen in the MCF-Adr cells (Figure 4B), further pointing to changes in MCF7-Adr that impaired the contribution of the ATF2 pathway to basal degree of programmed cell death.

Please replace the paragraph at page 46, lines 18-20 with the following:

Together, these findings establish that the expression of ATF2 peptides and in particular peptide II efficiently sensitizes melanoma and breast cancer cells to apoptosis induced by chemotherapeutic drugs, including MMC, Adriamycin® ~~adriamycin~~ + verapamil and UCN-01.

Please replace the paragraph at page 49, line 9 to page 50, line 1 with the following:

The present study has extended earlier observations in which ATF2 was identified as an important player in the melanoma cell's ability to undergo apoptosis. Four 50 amino acid peptides obtained from the amino-terminal domain of ATF2 were tested, of which two were selected for further characterization, on the basis of their pronounced effect on late-stage

Please replace the paragraph at page 52, line 29 to page 53, line 7 with the following:

Please replace the paragraph at page 59, lines 15-25 with the following:

Please replace the paragraph at page 63, lines 20-28 with the following:

GenePix 4000 4000® scanner (Union city, CA) and fluorescent data were collected using GenePix™ ~~GenePix~~ software.

Please replace the paragraph at page 63, line 29 to page 64, line 2, with the following:

Data Analysis. The axon image data for each microarray was uploaded to the NCI mAdb database for subsequent analysis using a variety of statistical web based tools (~~e.g.,~~ nciarray.nci.nih.gov). Gene clustering analysis was performed using the clustering algorithm and tree view software developed by Mike Eisen (Stanford, CA).